

8558  
DSSK



1977

was maintained. This fact probably explains the constancy of the RBC and Hc values compared with freshwater-transported animals where the decrease in blood Na<sup>+</sup> concentration is known to be associated with a decrease in blood osmotic pressure (Hattingh & Van Pletzen 1974). This eventually leads to haemolysis. Maintenance of a constant blood Na<sup>+</sup> concentration during transportation therefore seems to be crucial for these animals and they most probably do this by absorbing NaCl from their medium.

Finally, although it is apparent that addition of salt to the transportation water aids the fish, the decrease in blood pH on the second day is indicative that some stress is still experienced. The cause of this remains to be investigated.

REFERENCES

BOUCK, G. R. & BALL, R. C. 1966. Influence of capture methods on blood characteristics and mortality in the rainbow trout (*Salmo*

*gairdneri*). *Trans. Am. Fish. Soc.* 95: 170-175.  
FUJIYA, M. 1961. Use of electrophoretic serum separations in fish studies. *J. Wat. Pollut. Control Fed.* 33: 250-257.  
HATTINGH, J. & VAN PLETZEN, A. J. J. 1974. The influence of capture and transportation on some blood parameters of freshwater fish. *Comp. Biochem. Physiol.* 49A: 607-609.  
HATTINGH, J., FOURIE, F. LE R. & VAN VUREN, J. H. J. 1975. The transport of freshwater fish. *J. Fish. Biol.* 7: 447-449.  
MANN, K. H. 1965. Energy transformation by a population of fish in the River Thames. *J. anim. Ecol.* 34: 253-275.  
NARASIMHAN, P. V. & SUNDARARAY, B. I. 1971. Effects of stress on carbohydrate metabolism in the teleost *Notopterus notopterus* (Palles). *J. Fish. Biol.* 3: 441-452.  
WEDEMEYER, G. 1972. Some physiological consequences of handling stress in the juvenile coho salmon (*Oncorhynchus kisutch*) and steelhead trout (*Salmo gairdneri*). *J. Fish. Res. Bd Can.* 29: 1780-1783.

SAND SOLIDIFIED BY GEMSBOK URINE AS SELECTED BURROW SITES BY GERBILS

M. K. SEELY 1977

Desert Ecological Research Unit, Walvis Bay

Accepted: August, 1976.

*Gerbillus paeba* is widely distributed throughout the arid parts of southern Africa (Roberts 1951). Its range extends into the central Namib Desert where it is most often found on sandy areas of the central Namib plains or in the interdune valleys (Coetzee 1969). Its semi-permanent burrows are excavated in the sand either in the open (Laycock 1975) or, more often, close to dune-grass clumps (Coetzee 1969) and other vegetation. The sand

at the base of perennial vegetation clumps (e.g. *Acanthosicyos horrida*, *Stipagrostis sabulicola*, *Trianthema hereroensis*) is solidified by fog precipitation falling from the plants and by the presence of the roots which facilitate burrow construction. Distribution of gerbil burrows in dunes is thus limited by the occurrence of perennial vegetation or other suitable burrow sites in the more compact pebbly soils of the interdune valleys.

Unconsolidated sandy interdune valleys can be successfully colonized only under special conditions. One such opportunity is presented by the occurrence of sand patches which have been solidified by gemsbok (*Oryx gazella*) urine. The concentrated gemsbok urine lightly cements a block of sand averaging 0.025 m<sup>3</sup> (n = 20) in diameter which is more resistant to wind erosion and other disturbance than the surrounding loose

G. paeba / tytonis  
no date

is the study animal definitely paeba?  
The habitat suggests tytonis possibly.




FIGURE 1

Gerbil burrow at sand solidified by gemsbok urine which is situated in a sandy interdune valley.

u 2

sand. This provides a suitable location for the entrance to gerbil burrows which can then be extended into the more cohesive subsurface sands (Figure 1). Gemsbok occur widely and regularly throughout the dunes thus providing numerous suitable burrow sites.

In one sandy interdune valley 16 km south of Natab on the Kuiseb River in the Namib Desert Park a one-hectare plot was surveyed for gerbil holes and patches of sand solidified by gemsbok urine. Multiple entrances at one gemsbok urine patch were counted as one burrow. Of 134 gemsbok urine patches, 106 (79 per cent) were used for gerbil burrow entrances with from one to three entrances each. In this area there were 17 other gerbil burrow entrances. Thus of all burrow entrances in this area 86 per cent were situated at gemsbok urine patches. The surface area solidified by gemsbok urine totalled 13,4 m<sup>2</sup> or  $1,34 \times 10^{-5}$  per cent of the total area surveyed. Using the chi-square test a non-random distribution of burrow sites was demonstrated ( $p < 0,001$ ).

This valley supports a sparse cover of dried annual *Stipagrostis gonatostachys*, *Monsonia ignorata* and *Asthenatherum glaucum*, none of which is suitable for gerbil burrow entrances. Gerbils are highly mobile, crossing high dunes (Coetzee 1969) and are sometimes found far from their burrows (Laycock 1975). At such times they are subject to predation, primarily by jackals (*Canis mesomelas*). Access to escape

provided by foraging in the vicinity of burrow systems probably reduces the possibility of being preyed upon. Thus the advantage gained by a more even distribution of burrows relative to food sources is facilitated by the use of solidified gemsbok urine sand patches as burrow sites.

#### ACKNOWLEDGEMENTS

Mr P. K. Mostert and Miss A. Henderson are thanked for their assistance in the field. The continued financial support of the CSIR and Transvaal Museum is gratefully acknowledged. Permission to work in the Namib Desert Park was granted by the Division of Nature Conservation and Tourism, South West Africa.

#### REFERENCES

- COETZEE, C. G. 1969. The distribution of mammals in the Namib desert and adjoining inland escarpment. *Scient. Pap. Namib Des. Res. Stn*, 40: 23-36.
- LAYCOCK, P. A. 1975. A brief gerbil trapping survey in an interdune valley in the Namib Desert. *Madoqua* (2) 4: 95-97.
- ROBERTS, A. 1951. *The mammals of South Africa*. Johannesburg: Trustees 'The mammals of South Africa Book Fund.'

#### BABOONS AS DISPERSAL AGENTS FOR ACACIA CYCLOPS

CHRISTINE DAVIDGE

Zoology Department, University of Cape Town

Accepted: October, 1976.

Baboons, many birds and a few rodents in the south-west Cape include significant proportions of seeds of the Australian wattle (*Acacia cyclops* - 'rooikranz') in their diets (Middlemiss 1963, 1974). These animals may, therefore, act as dispersal agents for *A. cyclops* by distributing undigested seeds in their droppings. A one-year study (March 1975-February 1976) of a troop of chacma baboons, *Papio ursinus*, in the Cape of Good Hope Nature Reserve provided oppor-

tunities for gathering information on the part played by baboons in the spread of *A. cyclops* in the reserve.

Adult and juvenile baboons were observed eating *A. cyclops* seeds throughout the year. Greatest utilization of the seeds, however, occurred in April and May when, on average, an adult male baboon consumed daily about 19 000 *A. cyclops* seeds (Davidge 1976). Adult female and juvenile baboons appeared to take similar amounts of the seeds. The 85 baboons of the troop studied therefore consume large quantities of *A. cyclops* seeds each year.

Not all seeds ingested by baboons are cracked during mastication. Undamaged seeds are not digested during passage through the gut, but the seed tests are chemically and mechanically abraded. To determine the effect of passage through the gut on the rate of germination of *A. cyclops* seeds, one hundred intact seeds were collected from several dry baboon droppings found near Olifantsbos in the reserve. A further one hundred fully-developed seeds were collected from pods still attached to trees. These seeds were used as a control, since they were unlikely to have been mechanically abraded. The two sets of seeds ('ingested' and 'control') were arranged in rows on sandy soil 10 cm deep in a glass tank. The seeds were covered with a uniform layer of soil 5 cm deep; a surface partition separated the two sets of seeds. The soil was kept damp by frequent watering. The first 'ingested' seeds germinated seven days earlier than did the first 'control' seeds, and at a faster rate: 25 'ingested'

and five 'control' seeds had germinated after three weeks. Roughly equal numbers of 'control' and 'ingested' seeds (43 and 39 respectively) had, however, germinated 13 weeks after planting. It appears, therefore, that passage through the gut of a baboon does not significantly affect the amount of germination over three months, although the rate of germination was probably increased.

The baboons usually defaecated on top of rocks and on tarred roads, environments unfavourable for germination of seeds. Thirty baboon droppings, examined in February, each contained an average of 20 (range 1-200) undigested *A. cyclops* seeds. The undigested seeds appeared (from the remains of finely divided arils and tests also in the droppings) to be only a small proportion of all ingested seeds. Thus it seems unlikely that baboons are important agents in the dispersal of *A. cyclops* seeds in the Cape of Good Hope Nature Reserve.

#### REFERENCES

- DAVIDGE, C. 1976. Activity patterns of baboons *Papio ursinus* at Cape Point. M.Sc. Thesis, Zoology Department, University of Cape Town.
- MIDDLEMISS, E. 1963. Distribution of *Acacia cyclops* in the Cape Peninsula area by birds and other animals. *S. Afr. J. Sci.* 59: 419-420
- MIDDLEMISS, E. 1974. *The Rondevlei Bird Sanctuary*. Divisional Council Cape.

---

#### FOOD, BREEDING AND COAT COLOUR OF FERAL CATS ON DASSEN ISLAND

J. COOPER  
*FitzPatrick Institute,  
University of Cape Town*

Accepted: October, 1976.

Observations of food, breeding and coat colour of feral cats *Felis domesticus* were made from December 1971 to November 1972 on Dassen Island, South Africa (33°26'S/18°05'E). Five species of birds were recorded as prey, either by observation of cats feeding or by finding tooth